U.S. FISH AND WILDLIFE SERVICE SPECIES ASSESSMENT AND LISTING PRIORITY ASSIGNMENT FORM

SCIENTIFIC NAME: Pittosporum napaliense
COMMON NAME: Ho`awa
LEAD REGION: Region 1
INFORMATION CURRENT AS OF: June 2004
STATUS/ACTION
 Species assessment - determined species did not meet the definition of endangered or threatened under the Act and, therefore, was not elevated to Candidate status New candidate
X Continuing candidate
Non-petitioned
X Petitioned - Date petition received: May 11, 2004
_ 90-day positive - FR date:
X 12-month warranted but precluded - FR date: May 11, 2005
N Did the petition request a reclassification of a listed species?
FOR PETITIONED CANDIDATE SPECIES:
a. Is listing warranted (if yes, see summary of threats below)? <u>yes</u>b. To date, has publication of a proposal to list been precluded by other higher priority
listing actions? <u>yes</u>
c. If the answer to a. and b. is "yes", provide an explanation of why the action is
precluded. We find that the immediate issuance of a proposed rule and timely
promulgation of a final rule for this species has been, for the preceding 12 months, and
continues to be, precluded by higher priority listing actions. During the past 12 months,
most of our national listing budget has been consumed by work on various listing actions
to comply with court orders and court-approved settlement agreements, meeting statutory
deadlines for petition findings or listing determinations, emergency listing evaluations
and determinations and essential litigation-related, administrative, and program
management tasks. We will continue to monitor the status of this species as new
information becomes available. This review will determine if a change in status is
warranted, including the need to make prompt use of emergency listing procedures. For
information on listing actions taken over the past 12 months, see the discussion of
"Progress on Revising the Lists," in the current CNOR which can be viewed on our
Internet website (http://endangered.fws.gov).
Listing priority change
Former LP: New LP:
Date when the species first became a Candidate (as currently defined): 1990
Candidate removal: Former LP:
A – Taxon is more abundant or widespread than previously believed or not subject to
—— r

the degree of threats sufficient to warrant issuance of a proposed listing or
continuance of candidate status.
U – Taxon not subject to the degree of threats sufficient to warrant issuance of a proposed listing or continuance of candidate status due, in part or totally, to conservation efforts that remove or reduce the threats to the species.
F – Range is no longer a U.S. territory.
I – Insufficient information exists on biological vulnerability and threats to suppor
listing.
M – Taxon mistakenly included in past notice of review.
N – Taxon does not meet the Act's definition of "species."
X – Taxon believed to be extinct.

ANIMAL/PLANT GROUP AND FAMILY: Flowering plants, Pittosporaceae (Pittosporum family)

HISTORICAL STATES/TERRITORIES/COUNTRIES OF OCCURRENCE: Hawaii, island of Kauai

CURRENT STATES/ COUNTIES/TERRITORIES/COUNTRIES OF OCCURRENCE: Hawaii, island of Kauai

LAND OWNERSHIP: All populations occur on State lands.

LEAD REGION CONTACT: Paul Phifer, 503-872-2823, paul_phifer@fws.gov

LEAD FIELD OFFICE CONTACT: Pacific Islands Fish and Wildlife Office, Christa Russell, 808-792-9400, christa_russell@fws.gov

BIOLOGICAL INFORMATION:

<u>Species Description</u> *Pittosporum napaliense* is a small tree, 5 to 11 meters (m) (16 to 36 feet (ft)) tall with young parts and inflorescences covered in pale brown pubescence. The glabrous leaves are obovate with entire margins and slightly impressed veins on the upper surface. The cream colored, tubular flowers are unisexual and occur in groups of 8 to 20 in axillary or terminal inflorescences. Capsules are oblong-ovoid, 17 to 19 millimeters (0.06 to 0.07 inches) long, and glabrous at maturity (Wagner *et al.* 1999a).

<u>Taxonomy</u> *Pittosporum napaliense* was described by Sherff. This species is recognized as a distinct taxon in Wagner *et al.* (1999a) and Wagner and Herbst (2003), the most recently accepted Hawaiian plant taxonomy.

<u>Habitat</u> Typical habitat is *Pandanus* forest and mesic valleys at elevations between 150 and 550 m (490 to 1,800 ft) (Wagner *et al.* 1999a).

<u>Historical and Current Range/Current Status</u> *Pittosporum napaliense* is known from about six populations totaling several hundred individuals. This species is found on the eastern portion of

the Na Pali coast on the island of Kauai (Steve Perlman, National Tropical Botanical Garden, pers. comm. 1995).

THREATS:

A. The present or threatened destruction, modification, or curtailment of its habitat or range. This species is threatened by feral pigs (Sus scrofa) (S. Perlman, pers. comm. 1995). As early as 1778, European explorers introduced livestock, which became feral, increased in number and range, and caused significant changes to the natural environment of Hawaii. Past and present activities of introduced alien mammals are the primary factor altering and degrading vegetation and habitats on Kauai. The pig is originally native to Europe, northern Africa, Asia Minor, and Asia. European pigs, introduced to Hawaii by Captain James Cook in 1778, became feral and invaded forested areas, especially wet and mesic forests and dry areas at high elevations. They are currently present on Kauai and four other islands, and inhabit rain forests and grasslands. While rooting in the ground in search of the invertebrates and plant material they eat, feral pigs disturb and destroy vegetative cover, trample plants and seedlings, and threaten forest regeneration by damaging seeds and seedlings. They disturb soil and cause erosion, especially on slopes. Alien plant seeds are dispersed on their hooves and coats as well as through their digestive tracts, and the disturbed soil is fertilized by their feces, helping these plants to establish. Pigs are a major vector in the spread of many introduced plant species (Smith 1985; Stone 1985; Medeiros et al. 1986; Scott et al. 1986; Tomich 1986; Cuddihy and Stone 1990; Wagner et al. 1999a). No known conservation measures have been implemented to date to address this threat.

B. Overutilization for commercial, recreational, scientific, or educational purposes. None known.

C. <u>Disease or predation</u>.

None known.

D. The inadequacy of existing regulatory mechanisms.

Pigs are managed in Hawaii as game animals but may populate inaccessible areas where hunting is difficult, if not impossible, and therefore has little effect on their numbers (Hawaii Heritage Program 1990). Pig hunting is allowed on all islands either year-round or during certain months, depending on the area (Hawaii Department of Land and Natural Resources n.d.-a, n.d.-b, n.d.-c, n.d.-d). However, public hunting does not adequately control the number of pigs to eliminate this threat to this species. No other known conservation measures have been implemented to date to address this threat.

E. Other natural or manmade factors affecting its continued existence.

This species is threatened by alien plant species that adversely modify habitat (S. Perlman, pers. comm. 1995).

The original native flora of Hawaii consisted of about 1,400 species, nearly 90 percent of which were endemic. Of the total native and naturalized Hawaiian flora of 1,817 taxa, 47 percent were introduced from other parts of the world, and nearly 100 species have become pests (Smith 1985;

Wagner *et al.* 1999a). Several studies (Cuddihy and Stone 1990; Wood and Perlman 1997; Robichaux *et al.* 1998) indicate nonnative plant species may outcompete native plants similar to *Pittosporum napaliense*. Competition may be for space, light, water, or nutrients, or there may be a chemical inhibition of other plants (Smith 1985; Cuddihy and Stone 1990). In addition, nonnative pest plants found in habitat similar to that of this species have been shown to make the habitat less suitable for native species (Smathers and Gardner 1978; Smith 1985; Loope and Medeiros 1992; Medeiros *et al.* 1992; Ellshoff *et al.* 1995; Meyer and Florence 1996; Medeiros *et al.* 1997; Loope *et al.* 2004). In particular, alien pest plant species modify habitat by modifying availability of light, altering soil-water regimes, modifying nutrient cycling, or altering fire characteristics of native plant communities (Smith 1985; Cuddihy and Stone 1990; Vitousek *et al.* 1987). Because of demonstrated habitat modification and resource competition by nonnative plant species in habitat similar to habitat of *Pittosporum napaliense*, the Service believes nonnative plant species are a threat to *Pittosporum napaliense*. The remaining unmanaged populations of *Pittosporum napaliense* are still impacted by this threat.

CONSERVATION MEASURES PLANNED OR IMPLEMENTED None known.

SUMMARY OF THREATS

The major threats to this species include feral pigs and nonnative plant species, which are believed to be a major cause of the decline of this species throughout its range. No conservation efforts have been initiated to date.

LISTING PRIORITY

THREAT			
Magnitude	Immediacy	Taxonomy	Priority
High	Imminent Non-imminent	Monotypic genus Species Subspecies/population Monotypic genus Species Subspecies/population	1 2* 3 4 5 6
Moderate to Low	Imminent Non-imminent	Monotypic genus Species Subspecies/population Monotypic genus Species Subspecies/population	7 8 9 10 11 12

Rationale for listing priority number:

Magnitude:

This species is highly threatened by feral pigs that degrade and destroy habitat, and non-native

plants that compete for light and nutrients. Threats to the *Pandanus* forest and mesic valley habitat of *Pittosporum napaliense* and to individuals of this species occur throughout its range and are expected to continue or increase without their control or eradication. No conservation efforts have been initiated to date.

Imminence:

Threats to *Pittosporum napaliense* from feral pigs and non-native plants are considered imminent because they are ongoing.

<u>Yes</u> Have you promptly reviewed all of the information received regarding the species for the purpose of determining whether emergency listing is needed?

Is Emergency Listing Warranted? No. The species does not appear to be appropriate for emergency listing at this time because the immediacy of the threats is not so great as to imperil a significant proportion of the taxon within the time frame of the routine listing process. If it becomes apparent that the routine listing process is not sufficient to prevent large losses that may result in this species' extinction, then the emergency rule process for this species will be initiated. We will continue to monitor the status of *Pittosporum napaliense* as new information becomes available. This review will determine if a change in status is warranted, including the need to make prompt use of emergency listing procedures.

DESCRIPTION OF MONITORING:

The information in this form is based on the results of a meeting of 20 botanical experts held by the Center for Plant Conservation in December of 1995, and was updated by personal communication with Steve Perlman of the National Tropical Botanical Garden. We have incorporated additional information on this species from our files and the most recent supplement to the *Manual of the Flowering Plants of Hawaii* (Wagner and Herbst 2003). In 2004 the Pacific Islands office contacted the following species experts: Bob Hobdy, retired from Hawaii Division of Forestry and Wildlife; Joel Lau, Hawaii Natural Heritage Program; Art C. Medeiros III, U.S.G.S. Biological Resources Discipline; Hank Oppenheimer, resource manager for Maui Land and Pineapple Company; and Steve Perlman and Ken Wood, National Tropical Botanical Garden. No new information was provided in 2004. In 2005 we contacted the species experts listed below, but received no new information on this taxon.

The Hawaii Natural Heritage Program identified this species as critically imperiled (Hawaii Natural Heritage Program Database 2004). Based on the International Union for Conservation of Nature and Natural Resources Red Plant Data Book rarity categories, this species is recognized as Rare (could be considered at risk) by Wagner *et al.* (1999b)

Species experts were contacted but did not provide new information this year, no new literature was found, and no known entities are studying this species. However, it is highly likely that the previously reported threats continue to impact the species at the same or an increased level.

COORDINATION WITH STATES

In October 2004 we provided the Hawaii Division of Forestry and Wildlife with copies of our

most recent candidate assessments for their review and comment. Vickie Caraway, the State botanist, reviewed the information for this species and provided no additional information or corrections (V. Caraway, pers. comm. 2005).

LITERATURE CITED

List all experts contacted:

Name		Date	Place of Employment
1.	Joel Lau	June 28, 2005	Hawaii Natural Heritage Program
2.	Art Medeiros	June 28, 2005	U.S.G.S. Biological Resources Discipline
3.	Jim Jacobi	June 28, 2005	U.S.G.S. Biological Resources Discipline
4.	Rick Warshauer	June 28, 2005	U.S.G.S. Biological Resources Discipline
5.	Hank Oppenheimer	June 28, 2005	Maui Land and Pineapple Company
6.	Kapua Kawelo	June 28, 2005	U.S. Army
7.	Dave Lorence	June 28, 2005	National Tropical Botanical Garden
8.	Steve Perlman	June 28, 2005	National Tropical Botanical Garden
9.	Ken Wood	June 28, 2005	National Tropical Botanical Garden
10.	Marie Bruegmann	July 13, 2005	U.S. Fish and Wildlife Service
11.	Vickie Caraway	June 14, 2005	Hawaii Division of Forestry and Wildlife

List all databases searched:

Name Date

1. Hawaii Natural Heritage Program 2004

Other resources utilized:

- Center for Biological Diversity, Dr. Jane Goodall, Dr. E.O. Wilson, Dr. Paul Ehrlich, Dr. John Terborgh, Dr. Niles Eldridge, Dr. Thomas Eisner, Dr. Robert Hass, Barbara Kingsolver, Charles Bowden, Martin Sheen, the Xerces Society, and the Biodiversity Conservation Alliance. 2004. Hawaiian Plants: petitions to list as federally endangered species. May 4, 2004.
- Cuddihy, L.W., and C.P. Stone. 1990. Alteration of native Hawaiian vegetation; effects of humans, their activities and introductions. Coop. Natl. Park Resources Stud. Unit, Hawaii. 138 pp.
- Ellshoff, Z.E., D.E. Gardner, C. Wikler, and C.W. Smith. 1995. Annotated bibliography of the genus *Psidium*, with emphasis on *P. cattleianum* (strawberry guava) and *P. guajava* (common guava), forest weeds in Hawai`i. Cooperative National Park Resources Studies Unit, University of Hawaii. Technical Report 95.
- Hawaii, Department of Land and Natural Resources. N.d.-a. Summary of Title 13, Chapter 123, Game mammal hunting rules, island of Oahu. Division of Forestry and Wildlife, Honolulu. 2 pp.
- Hawaii, Department of Land and Natural Resources. N.d.-b. Summary of Title 13, Chapter 123, Game mammal hunting rules, island of Molokai. Division of Forestry and Wildlife, Honolulu. 2 pp.
- Hawaii, Department of Land and Natural Resources. N.d.-c. Summary of Title 13, Chapter 123, Game mammal hunting rules, island of Maui. Division of Forestry and Wildlife, Honolulu. 2 pp.

- Hawaii, Department of Land and Natural Resources. N.d.-d. Summary of Title 13, Chapter 123, Game mammal hunting rules, island of Kauai. Division of Forestry and Wildlife, Honolulu.
- Loope, L.L. and A.C. Medeiros. 1992. A new and invasive grass on Maui. Newsletter of the Hawaiian Botanical Society 31: 7-8.
- Loope, L., F. Starr and K. Starr. 2004. Management and research for protecting endangered Hawaiian plant species from displacement by invasive plants on Maui, Hawaii. Weed Technology 18: 1472-1474.
- Medeiros, A.C., L.L. Loope, P. Conant and S. McElvaney. 1997. Status, ecology, and management of the invasive plant, *Miconia calvescens* DC (Melastomataceae) in the Hawaiian Islands. Bishop Mus. Occas. Pap. 48: 23-36.
- Medeiros, A.C., L.L. Loope, T. Flynn, S.J. Anderson, L.W. Cuddihy, and K.A. Wilson. 1992. Notes on the status of an invasive Australian tree fern (*Cyathea cooperi*) in Hawaiian rain forests. American Fern Journal 82: 27-33.
- Medeiros, A.C., Jr., L.L. Loope, and R.A. Holt. 1986. Status of native flowering plant species on the south slope of Haleakala, East Maui, Hawaii. Coop. Natl. Park Resources Stud. Unit, Hawaii, Techn. Rept. 59:1-230.
- Meyer, J.-Y. and J. Florence. 1996. Tahiti's native flora endangered by the invasion of *Miconia calvescens* D.C. (Melastomataceae). Journal of Biogeography 23: 775-781.
- Robichaux, R., J. Canfield, F. R. Warshauer, L. Perry, M. Bruegmann, and G. Carr. 1998. Adaptive Radiation. Endangered Species Bulletin. November/December.
- Scott, J.M., S. Mountainspring, F.L. Ramsey, and C.B. Kepler. 1986. Forest bird communities of the Hawaiian Islands: Their dynamics, ecology, and conservation. Studies in Avian Biology 9:1-429. Cooper Ornithological Society, Los Angeles.
- Smathers, G.A. and D.E. Gardner. 1978. Stand analysis of an invading firetree (*Myrica faya* Aiton) population, Hawai`i. Proceeding of the Second Conference on Natural Science, Hawaii Volcanoes National Park, pp. 274-288.
- Smith, C.W. 1985. Impact of alien plants on Hawai`i's native biota: <u>in</u> Stone, C.P., and J.M. Scott (eds.), Hawai`i's terrestrial ecosystems: preservation and management. Coop. Natl. Park Resources Stud. Unit, Univ. Hawaii, Honolulu, pp. 180-250.
- Stone, C.P. 1985. Alien animals in Hawai`i's native ecosystems: toward controlling the adverse effects of introduced vertebrates: <u>in</u> Stone, C.P., and J.M. Scott (eds.), Hawai'i's terrestrial ecosystems: preservation and management. Coop. Natl. Park Resources Stud. Unit, Univ. Hawaii, Honolulu, pp. 251-297.
- Tomich, P.Q. 1986. Mammals in Hawai`i; a synopsis and notational bibliography. Bishop Museum Press, Honolulu. 375 pp.
- Vitousek, P.M., C.M. D'Antonio, L.L. Loope, M. Rejnanek, and R. Westerbrooks. 1997. Introduced species: a significant component of human-caused global change. New Zealand Journal of Ecology 21(1): 1-16.
- Wagner, W.L., D.R. Herbst, and S.H. Sohmer. 1999a. Manual of the Flowering Plants of Hawai`i, Bishop Mus. Spec. Publ. 97:1-1918. University of Hawaii Press and Bishop Museum Press, Honolulu.
- Wagner, W.L., M.M. Bruegmann, and J.Q.C. Lau. 1999b. Hawaiian vascular plants at risk: 1999. Bishop Mus. Occas. Pap. 60: 1-58.
- Wagner, W.L. and D.R. Herbst. 2003. Electronic supplement to the manual of flowering plants

of Hawai'i, version 3.1. December 12, 2003. Available from the Internet. URL: http://rathbun.si.edu/botany/pacificislandbiodiversity/hawaiianflora/supplement.htm. Wenkam, R. 1969. Kauai and the park country of Hawaii. Sierra Club, San Francisco. 160 pp. Wood, K.R. and S. Perlman. 1997. Maui 14 plant survey final report. Submitted by National Tropical Botanical Garden, October, 1997.

APPROVAL/CONCURRENCE: Lead Regions must obtain written concurrence from all other Regions within the range of the species before recommending changes to the candidate list, including listing priority changes; the Regional Director must approve all such recommendations. The Director must concur on all 12-month petition findings, additions of species to the candidate list, removal of candidate species, and listing priority changes.

Approve:	Regional Director, Fish and Wildlife	Te Service Date
	Mauhau Smoge	
Concur:	Director, Fish and Wildlife Service	<u>August 23, 2006</u> Date
Do not concur	:	Date
	l review: <u>September 20, 2005</u> : <u>Marie M. Bruegmann, Pacific Island</u> Plant Recovery Coordinator	<u>ds FWO</u>
Comments: PIFWO Revie Reviewed by:	w Christa Russell Plant Conservation Program Leader	Date: September 27, 2005
	Gina Shultz Assistant Field Supervisor, Endangered Species	Date: October 14, 2005
	Patrick Leonard Field Supervisor	Date: October 14, 2005